

JOURNAL OF ALGEBRA 122, 64–80 (1989)

2-Cocycles on the Algebra of Differential Operators*

WANG-LAI LI[†]

*Institute of Systems Science, Academia Sinica,
Beijing, People's Republic of China*

Communicated by Nathan Jacobson

Received June 10, 1987

1. INTRODUCTION

The central extension method is often used in the theory of structures and representations of Kac–Moody algebras. By central extension, we can construct many infinite-dimensional Lie algebras, such as affine Lie algebras, infinite-dimensional Heisenberg algebras, and Virasoro algebras, which have a profound mathematical and physical background [1, 5, 7, 8]. We can describe the structures and some of the representations of these Lie algebras by using this method [4, 9].

If \mathfrak{G} is a Lie algebra, it is well known that all 1-dimensional central extensions of \mathfrak{G} are determined by the 2-cocycle group of \mathfrak{G} . If a 2-cocycle is induced by a linear function on \mathfrak{G} , this 2-cocycle is called trivial; in this case, the 1-dimensional central extension is also trivial, i.e., it is isomorphic to $\mathfrak{G} \oplus \mathbb{C}c$ as a direct sum of ideals. If we want to describe all 1-dimensional central extensions of \mathfrak{G} , it is necessary to determine all non-trivial 2-cocycles or the second cohomology group of \mathfrak{G} [3].

Let $\mathbb{C}[t, t^{-1}]$ be the Laurent polynomial ring over a complex field \mathbb{C} , $\alpha_1(t^m, t^n) = m\delta_{m, -n}$, $m, n \in \mathbb{Z}$. We can construct a 1-dimensional central extension of $\mathbb{C}[t, t^{-1}]$ by using α_1 , and the Lie bracket on $\mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$ is defined by

$$\begin{aligned} [t^m, t^n] &= \alpha_1(t^m, t^n) c, \\ [c, t^n] &= 0, \quad [c, c] = 0, \quad m, n \in \mathbb{Z}. \end{aligned}$$

* Project supported by the Science Fund of the Chinese Academy of Sciences.

[†] Current address: Department of Mathematics, Beijing Normal University, Beijing, People's Republic of China.

This is an infinite-dimensional Lie algebra. The subalgebra

$$\bigoplus_{\substack{n \in \mathbb{Z} \\ n \neq 0}} \mathbb{C} t^n \oplus \mathbb{C} c$$

is an infinite-dimensional Heisenberg algebra.

Denote the differential operator d/dt on $\mathbb{C}[t, t^{-1}]$ by D . Let $\mathfrak{L} = \{P(t) D \mid P(t) \in \mathbb{C}[t, t^{-1}]\}$ be the algebra of all derivations on $\mathbb{C}[t, t^{-1}]$,

$$\alpha_2(t^{i+1}D, t^{j+1}D) = \frac{1}{12}(i^3 - i) \delta_{i, -j}, \quad i, j \in \mathbb{Z}.$$

We can construct a 1-dimensional central extension of \mathfrak{L} by using α_2 , and the Lie bracket on $\mathfrak{L} \oplus \mathbb{C}c$ is defined by

$$\begin{aligned} [t^{i+1}D, t^{j+1}D] &= (j-i) t^{i+j+1}D + \alpha_2(t^{i+1}D, t^{j+1}D) c, \\ [c, t^{i+1}D] &= 0, \quad [c, c] = 0, \quad i, j \in \mathbb{Z}. \end{aligned}$$

This is an infinite-dimensional Lie algebra and is called a Virasoro algebra.

Let $\mathfrak{a} = \mathbb{C}[t, t^{-1}, D]$ be the algebra of differential operators on $\mathbb{C}[t, t^{-1}]$,

$$\alpha_3(t^{l+m}D^l, t^{l'+m'}D^{l'}) = \delta_{m, -m'}(-1)^l l! l'! \binom{m+l}{l+l'+1},$$

$l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}$. We can construct a 1-dimensional central extension of \mathfrak{a} by using α_3 , and the Lie bracket on $\mathfrak{a} \oplus \mathbb{C}c$ is defined by

$$\begin{aligned} [t^{l+m}D^l, t^{l'+m'}D^{l'}] &= \sum_{i=0}^l \binom{l}{i} t^{l+m}(D^{l-i}(t^{l'+m'})) D^{l'+i} \\ &\quad - \sum_{j=0}^{l'} \binom{l'}{j} t^{l'+m'}(D^{l'-j}(t^{l+m})) D^{l+j} \\ &\quad + \alpha_3(t^{l+m}D^l, t^{l'+m'}D^{l'}) c, \\ [c, t^{l+m}D^l] &= 0, \quad [c, c] = 0, \quad l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}. \end{aligned}$$

Generally, let us consider the algebra of differential operators $\mathfrak{A} = \mathfrak{a} \otimes gl_n(\mathbb{C})$ on $\mathbb{C}[t, t^{-1}] \otimes gl_n(\mathbb{C})$, and set

$$\alpha_4(t^{l+m}D^l \otimes A, t^{l'+m'}D^{l'} \otimes A') = \delta_{m, -m'}(-1)^l l! l'! \binom{m+l}{l+l'+1} \text{tr } AA',$$

$l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}, A, A' \in gl_n(\mathbb{C})$. We can construct a 1-dimensional central extension of \mathfrak{A} by using α_4 , and the Lie bracket is defined by

$$\begin{aligned} & [t^{l+m}D^l \otimes A, t^{l'+m'}D^{l'} \otimes A'] \\ &= \sum_{i=0}^l \binom{l}{i} t^{l+m}(D^{l-i}(t^{l'+m'})) D^{l'+i} \otimes AA' \\ &\quad - \sum_{j=0}^{l'} \binom{l'}{j} t^{l'+m'}(D^{l'-j}(t^{l+m})) D^{l+j} \otimes A'A \\ &\quad + \alpha_4(t^{l+m}D^l \otimes A, t^{l'+m'}D^{l'} \otimes A') c, \end{aligned}$$

$l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}, A, A' \in gl_n(\mathbb{C})$.

It is easy to see that a non-trivial 1-dimensional central extension of $\mathbb{C}[t, t^{-1}]$ is not unique. All non-zero 2-cocycles on $\mathbb{C}[t, t^{-1}]$ are non-trivial and the second cohomology group is an infinite-dimensional vector space. It is known [2] that every 2-cocycle on \mathfrak{Q} is equivalent to a multiple of α_2 . In the lectures at the Graduate School, Academia Sinica, in 1985, V. G. Kac proposed the following open problem: "Is every 2-cocycle on \mathfrak{Q} equivalent to a multiple of α_3 ?" Later, Kac proposed a more general problem [6]: "Is every 2-cocycle on \mathfrak{A} equivalent to a multiple of α_4 ?"

In this paper, we give these problems a positive answer.

2. BASIC CONCEPTS AND MAIN RESULTS

We assume that the underlying field is complex field \mathbb{C} , though all results hold for all fields of characteristic 0.

Let \mathfrak{G} be a Lie algebra. A 2-cocycle on \mathfrak{G} is a bilinear \mathbb{C} -valued form ψ satisfying the following conditions:

- (i) $\psi(a, b) = -\psi(b, a)$,
- (ii) $\psi([a, b], c) + \psi([b, c], a) + \psi([c, a], b) = 0$,

for $a, b, c \in \mathfrak{G}$.

If f is a linear function on \mathfrak{G} , we define

$$\alpha_f(x, y) = f([x, y])$$

for $x, y \in \mathfrak{G}$. One easily checks that this is a 2-cocycle. This 2-cocycle is called a trivial 2-cocycle. A 2-cocycle φ is equivalent to a 2-cocycle ψ , if $\varphi - \psi$ is trivial.

Given a 2-cocycle α on \mathfrak{G} , we can construct a central extension of \mathfrak{G} . If the Lie bracket on \mathfrak{G} is $[\cdot, \cdot]$, we define a Lie bracket $[\cdot, \cdot]_0$ on $\mathfrak{G} \oplus \mathbb{C}c$ as

$$[x + \lambda c, y + \mu c]_0 = [x, y] + \alpha(x, y) c, \quad x, y \in \mathfrak{G}, \lambda, \mu \in \mathbb{C}.$$

It is easy to check that $\mathfrak{G} \oplus \mathbb{C}c$ is a Lie algebra with this bracket and every 1-dimensional central extension of \mathfrak{G} can be obtained in this way. Denote this central extension of \mathfrak{G} by $\mathfrak{G}(\alpha)$.

Let γ be a trivial 2-cocycle induced by f and let α be a 2-cocycle; then

$$\begin{aligned} x &\mapsto x + f(x) c, \\ c &\mapsto c, \end{aligned}$$

$x \in \mathfrak{G}$, gives an isomorphism from $\mathfrak{G}(\alpha)$ to $\mathfrak{G}(\alpha + \gamma)$.

Let $\mathbb{C}[t, t^{-1}]$ be the Laurent polynomial ring over \mathbb{C} , $D = d/dt$ the differential operator on $\mathbb{C}[t, t^{-1}]$, and $\mathfrak{a} = \mathbb{C}[t, t^{-1}, D]$ the algebra of differential operators on $\mathbb{C}[t, t^{-1}]$. \mathfrak{a} has a basis $\{t^{l+m}D^l \mid l \geq 0, m \in \mathbb{Z}\}$, and

$$\begin{aligned} [t^{l+m}D^l, t^{l'+m'}D^{l'}] &= (t^{l+m}D^l) \circ (t^{l'+m'}D^{l'}) - (t^{l'+m'}D^{l'}) \circ (t^{l+m}D^l) \\ &= \sum_{i=0}^l \binom{l}{i} t^{l+m} (D^{l-i} (t^{l'+m'})) D^{l+i} \\ &\quad - \sum_{j=0}^{l'} \binom{l'}{j} t^{l'+m'} (D^{l'-j} (t^{l+m})) D^{l'+j}. \end{aligned}$$

In particular,

$$[t^{l+m}D^l, D] = -(l+m) t^{l+m-1}D^l, \quad (2.1)$$

$$[t^{-1}D^l, tD] = (l+1) t^{-1}D^l. \quad (2.2)$$

From (2.1), (2.2) we have

$$[\mathfrak{a}, \mathfrak{a}] = \mathfrak{a}. \quad (2.3)$$

Let $gl_n(\mathbb{C})$ be an $n \times n$ matrix algebra over \mathbb{C} , and $\mathfrak{A} = \mathfrak{a} \otimes gl_n(\mathbb{C})$ the algebra of differential operators on $\mathbb{C}[t, t^{-1}] \otimes gl_n(\mathbb{C})$.

Define a \mathbb{C} -valued bilinear form α on \mathfrak{a} :

$$\alpha(t^{l+m}D^l, t^{l'+m'}D^{l'}) = \delta_{m, -m'} (-1)^l l! l'! \binom{m+l}{l+l'+1}, \quad (2.4)$$

$l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}$. From [5], α is a 2-cocycle on \mathfrak{a} .

Define a \mathbb{C} -valued bilinear form ψ on \mathfrak{A} :

$$\begin{aligned} \psi(t^{l+m}D^l \otimes A, t^{l'+m'}D^{l'} \otimes A') \\ = \delta_{m, -m'} (-1)^l l! l'! \binom{m+l}{l+l'+1} \text{tr } AA', \end{aligned} \quad (2.5)$$

$l \geq 0, l' \geq 0, m, m' \in \mathbb{Z}, A, A' \in gl_n(\mathbb{C})$. From [8], ψ is a 2-cocycle on \mathfrak{A} . V. G. Kac proposed the following problems:

PROBLEM 1. Is every 2-cocycle on \mathfrak{a} equivalent to a multiple of α ?

PROBLEM 2. Is every 2-cocycle on \mathfrak{A} equivalent to a multiple of ψ ?

We give these problems a positive answer.

THEOREM 2.1. Every 2-cocycle on \mathfrak{a} is equivalent to a multiple of α .

THEOREM 2.2. Every 2-cocycle on \mathfrak{A} is equivalent to a multiple of ψ .

3. PROOF OF THEOREM 2.1

Now we can solve Kac's first problem.

From (2.4) we have

$$\begin{aligned} \alpha(t^{l+m}D^l, t^{l'+m'}D^{l'}) &= 0 \\ \Leftrightarrow m \neq -m' \text{ or } m = -m', \quad 0 \leq m+l \leq l+l'. \end{aligned} \quad (3.1)$$

Let β be a 2-cocycle on \mathfrak{a} . We define a linear function f_β on \mathfrak{a} :

$$\begin{aligned} f_\beta(t^{l+m-1}D^l) &= -\frac{1}{l+m} \beta(t^{l+m}D^l, D), \quad l+m \neq 0, \\ f_\beta(t^{-1}D^l) &= \frac{1}{l+1} \beta(t^{-1}D^l, tD). \end{aligned}$$

Then $\beta_1 = \beta - \alpha_{f_\beta}$ is a 2-cocycle on \mathfrak{a} and equivalent to β . From (2.1), (2.2) we have

$$\beta_1(t^{l+m}D^l, D) = 0, \quad l+m \neq 0, \quad (3.2)$$

$$\beta_1(t^{-1}D^l, tD) = 0. \quad (3.3)$$

LEMMA 3.1. $\beta_1(t^{l+m}D^l, D) = 0, l \geq 0, m \in \mathbb{Z}$.

Proof. From (3.2) it is sufficient to consider the case $l+m=0$. Since

$$\begin{aligned} [D^l, tD] &= D^l \circ (tD) - tD^{l+1} = lD^l, \\ \beta_1(D^l, D) &= \beta_1(D^l, [D, tD]) \\ &= \beta_1([D^l, D], tD) + \beta_1(D, [D^l, tD]) \\ &= -l\beta_1(D^l, D), \end{aligned}$$

this implies

$$\beta_1(D', D) = 0.$$

LEMMA 3.2. $\beta_1(1, a) = 0$.

Proof.

$$\begin{aligned}\beta_1(1, a) &= \beta_1(1, [a, a]) \\ &= \beta_1([1, a], a) + \beta_1(a, [1, a]) \\ &= 0.\end{aligned}$$

PROPOSITION 3.3. *If $l + m \neq -1$, then*

$$\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) = -\frac{l'+m'}{l+m+1} \beta_1(t^{l+m+1}D^l, t^{l'+m'-1}D^{l'}).$$

Proof. $[D, t^s D^u] = D \circ (t^s D^u) - t^s D^{u+1} = s t^{s-1} D^u$. If $l + m \neq -1$, by Lemma 3.1 we have

$$\begin{aligned}\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) &= \beta_1\left(\frac{1}{l+m+1} [D, t^{l+m+1}D^l], t^{l'+m'}D^{l'}\right) \\ &= \frac{1}{l+m+1} \beta_1([D, t^{l'+m'}D^{l'}], t^{l+m+1}D^l) \\ &\quad + \frac{1}{l+m+1} \beta_1(D, [t^{l+m+1}D^l, t^{l'+m'}D^{l'}]) \\ &= -\frac{l'+m'}{l+m+1} \beta_1(t^{l+m+1}D^l, t^{l'+m'-1}D^{l'}).\end{aligned}$$

LEMMA 3.4. $\beta_1(t^{l+m}D^l, tD) = 0, l \geq 0, m \in \mathbb{Z}$.

Proof. From (3.3) it is sufficient to prove the lemma for the case $l + m \neq -1$. In this case, by Proposition 3.3 we have

$$\beta_1(t^{l+m}D^l, tD) = -\frac{1}{l+m+1} \beta_1(t^{l+m+1}D^l, D) = 0.$$

PROPOSITION 3.5. *If $m \neq -m'$ or $m = -m', 0 \leq m + l \leq l + l'$, then*

$$\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) = 0$$

(cf. (3.1)).

Proof. $[tD, t^{l+m}D^l] = mt^{l+m}D^l$.

If $m \neq 0$, we have

$$\begin{aligned}\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) &= \beta_1\left(\frac{1}{m}[tD, t^{l+m}D^l], t^{l'+m'}D^{l'}\right) \\ &= \frac{1}{m}\beta_1([tD, t^{l'+m'}D^{l'}], t^{l+m}D^l) \quad (\text{Lemma 3.4}) \\ &= -\frac{m'}{m}\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}).\end{aligned}$$

Thus, if $m \neq 0$, $m \neq -m'$, we have

$$\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) = 0.$$

If $m = 0$, $m' \neq 0$, by Proposition 3.3 we have

$$\beta_1(t^lD^l, t^{l'+m'}D^{l'}) = -\frac{l'+m'}{l+1}\beta_1(t^{l+1}D^l, t^{l'+m'-1}D^{l'}) = 0.$$

Hence, we always have

$$\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) = 0, \quad \text{if } m \neq -m'.$$

Now we assume $m = -m'$, $0 \leq m+l \leq l+l'$, then

$$l' + m' = l' - m \geq 0.$$

By Proposition 3.3,

$$\begin{aligned}\beta_1(t^{l+m}D^l, t^{l'-m}D^{l'}) &= \frac{-(l'-m)}{l+m+1} \cdot \frac{-(l'-m-1)}{l+m+2} \cdots \frac{0}{l+l'+1} \beta_1(t^{l+l'+1}D^l, t^{-1}D^{l'}) \\ &= 0.\end{aligned}$$

This completes the proof.

LEMMA 3.6.

$$(i) \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} \beta_1(t^{-1}, t^2D) \\ \beta_1(t^{-1}D, t^2) \end{pmatrix} = \begin{pmatrix} 2\beta_1(t, t^{-1}) \\ 3\beta_1(t, t^{-1}) \end{pmatrix}, \quad (3.4)$$

(ii) if $k \geq 2$,

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & -1 & 1 \\ 0 & 0 & \cdots & -1 & 0 & 1 \\ & \cdots & & \cdots & \cdots & \\ -1 & & 0 & \cdots & 0 & 0 & 1 \\ 1 & k \binom{2k-1}{k+1} & \cdots & 0 & 0 & 0 \end{pmatrix} \times \begin{pmatrix} \beta_1(t^{-1}, t^{k+1}D^k) \\ \beta_1(t^{-1}D, t^{k+1}D^{k-1}) \\ \vdots \\ \beta_1(t^{-1}D^{k-1}, t^{k+1}D) \\ \beta_1(t^{-1}D^k, t^{k+1}) \end{pmatrix} = \begin{pmatrix} b_0 \\ b_1 \\ \vdots \\ b_{k-1} \\ b_k \end{pmatrix}. \quad (3.5)$$

where

$$\begin{aligned} b_u &= \frac{1}{(k-u+1)(k+2)} \left(\sum_{j=0}^{k-u-1} \binom{k-u-1}{j} \right) \\ &\quad \times (-1) \cdots (-1-k+u+j) \beta_1(t^{-k+u+j-2}D^{u+j}, t^{k+2}) \\ &\quad - \sum_{i=0}^{k-u-1} \binom{k-u-1}{i} (k+2) \cdots (u+i+2) \beta_1(t^{u+i+1}D^i, t^{-1}D^u), \end{aligned}$$

$u=0, \dots, k-1$,

$$\begin{aligned} b_k &= \frac{1}{k-1} \left(\sum_{i=0}^{k-2} \binom{k}{i} \right) (k-1) \cdots i \beta_1(t^{k+i}D^i, t^{-k+1}D) \\ &\quad - \sum_{j=0}^{k-2} \binom{k}{j} (-k+1) \cdots (-2k+2+j) \beta_1(t^{-k+2+j}D^{j+1}, t^{k-1}). \end{aligned}$$

Proof. $[D^{l+1}, t^{l+l'+2}] = \sum_{i=0}^l \binom{l+1}{i} (l+l'+2) \cdots (l'+2+i) t^{l'+i+1} D^i$.

We have

$$\begin{aligned} &\beta_1([D^{l+1}, t^{l+l'+2}], t^{-1}D^{l'}) \\ &= \sum_{i=0}^l \binom{l+1}{i} (l+l'+2) \cdots (l'+2+i) \beta_1(t^{l'+i+1}D^i, t^{-1}D^{l'}). \end{aligned}$$

On the other hand,

$$\begin{aligned}
& \beta_1([D^{l+1}, t^{l+l'+2}], t^{-1}D^r) \\
&= \beta_1([D^{l+1}, t^{-1}D^r], t^{l+l'+2}) + \beta_1(D^{l+1}, [t^{l+l'+2}, t^{-1}D^r]) \\
&= \beta_1\left(\sum_{j=0}^l \binom{l+1}{j} (-1) \cdots (-1 - (l+1-j) + 1) \right. \\
&\quad \left. \times t^{-1-(l+1-j)} D^{r+j}, t^{l+l'+2}\right) \\
&= \sum_{j=0}^l \binom{l+1}{j} (-1) \cdots (-1 - l + j) \\
&\quad \times \beta_1(t^{-l+j-2} D^{r+j}, t^{l+l'+2}) \quad (\text{by Proposition 3.5}).
\end{aligned}$$

This implies

$$\begin{aligned}
& \sum_{i=0}^l \binom{l+1}{i} (l+l'+2) \cdots (l'+2+i) \beta_1(t^{l'+i+1} D^l, t^{-1}D^r) \\
&= \sum_{j=0}^l \binom{l+1}{j} (-1) \cdots (-1 - l + j) \beta_1(t^{-l+j-2} D^{r+j}, t^{l+l'+2}). \quad (3.6)
\end{aligned}$$

By Proposition 3.3,

$$\beta_1(t^{-2} D^{l+r}, t^{l+l'+2}) = (l+l'+2) \beta_1(t^{-1} D^{l+r}, t^{l+l'+1}).$$

Thus (3.6) can be rewritten as

$$\begin{aligned}
& -\beta_1(t^{-1} D^r, t^{l+l'+1} D^l) + \beta_1(t^{-1} D^{l+r}, t^{l+l'+1}) \\
&= \frac{1}{(l+1)(l+l'+2)} \left(\sum_{j=0}^{l-1} \binom{l+1}{j} (-1) \cdots (-1 - l + j) \right. \\
&\quad \left. \times \beta_1(t^{-l+j-2} D^{r+j}, t^{l+l'+2}) \right. \\
&\quad \left. - \sum_{i=0}^{l-1} \binom{l+1}{i} (l+l'+2) \cdots (l'+2+i) \beta_1(t^{l'+i+1} D^l, t^{-1}D^r) \right). \quad (3.7)
\end{aligned}$$

Setting $l' = 0, l = 1$ and using Proposition 3.3, we obtain

$$-\beta_1(t^{-1}, t^2 D) + \beta_1(t^{-1} D, t^2) = 2\beta_1(t^{-1}, t). \quad (3.8)$$

Since

$$\begin{aligned}
& \beta_1([tD^2, t^2], t^{-1}) = \beta_1(4t^2 D + 2t, t^{-1}) \\
&= -4\beta_1(t^{-1}, t^2 D) + 2\beta_1(t, t^{-1}), \\
& \beta_1([tD^2, t^2], t^{-1}) = \beta_1([tD^2, t^{-1}], t^2) + \beta_1(tD^2, [t^2, t^{-1}]) \\
&= \beta_1(-2t^{-1} D + 2t^{-2}, t^2) \\
&= -2\beta_1(t^{-1} D, t^2) + 2\beta_1(t^{-2}, t^2).
\end{aligned}$$

By Proposition 3.3, $\beta_1(t^{-2}, t^2) = 2\beta_1(t^{-1}, t)$, hence,

$$2\beta_1(t^{-1}, t^2D) - \beta_1(t^{-1}D, t^2) = 3\beta_1(t, t^{-1}). \quad (3.9)$$

From (3.8), (3.9), we have (3.4).

For $k \geq 2$,

$$\begin{aligned} & \beta_1([t^{k+1}D^k, t^{k-1}], t^{-k+1}D) \\ &= \sum_{j=0}^{k-1} \binom{k}{j} (k-1) \cdots j \beta_1(t^{k+j}D^j, t^{-k+1}D). \end{aligned}$$

On the other hand,

$$\begin{aligned} & \beta_1([t^{k+1}D^k, t^{k-1}], t^{-k+1}D) \\ &= \beta_1([t^{k+1}D^k, t^{-k+1}D], t^{k-1}) + \beta_1(t^{k+1}D^k, [t^{k-1}, t^{-k+1}D]) \\ &= \beta_1\left(\sum_{i=0}^{k-1} \binom{k}{i} (-k+1) \cdots (-2k+2+i) t^{-k+2+i}D^{i+1}, t^{k-1}\right) \\ &\quad - (k+1) \beta_1(tD^k, t^{k-1}) - (k-1) \beta_1(t^{k+1}D^k, t^{-1}) \\ &= k(-k+1) \beta_1(tD^k, t^{k-1}) - (k+1) \beta_1(tD^k, t^{k-1}) + (k-1) \\ &\quad \times \beta_1(t^{-1}, t^{k+1}D^k) \\ &\quad + \sum_{i=0}^{k-2} \binom{k}{i} (-k+1) \cdots (-2k+2+i) \beta_1(t^{-k+2+i}D^{i+1}, t^{k-1}) \\ &= (k-1) \beta_1(t^{-1}, t^{k+1}D^k) \\ &\quad + \sum_{i=0}^{k-2} \binom{k}{i} (-k+1) \cdots (-2k+2+i) \beta_1(t^{-k+2+i}D^{i+1}, t^{k-1}) \\ &\hspace{15em} \text{(by Proposition 3.5).} \end{aligned}$$

We have

$$\begin{aligned} & (k-1) \beta_1(t^{-1}, t^{k+1}D^k) - k(k-1) \beta_1(t^{2k-1}D^{k-1}, t^{-k+1}D) \\ &= \sum_{j=0}^{k-2} \binom{k}{j} (k-1) \cdots j \beta_1(t^{k+j}D^j, t^{-k+1}D) \\ &\quad - \sum_{i=0}^{k-2} \binom{k}{i} (-k+1) \cdots (-2k+2+i) \beta_1(t^{-k+2+i}D^{i+1}, t^{k-1}). \end{aligned}$$

By Proposition 3.3,

$$\beta_1(t^{2k-1}D^{k-1}, t^{-k+1}D) = -\binom{2k-1}{k+1} \beta_1(t^{-1}D, t^{k+1}D^{k-1}).$$

We have

$$\begin{aligned}
 & \beta_1(t^{-1}, t^{k+1}D^k) + k \binom{2k-1}{k+1} \beta_1(t^{-1}D, t^{k+1}D^{k-1}) \\
 &= \frac{1}{k-1} \left(\sum_{j=0}^{k-2} \binom{k}{j} (k-1) \cdots j \beta_1(t^{k+j}D^j, t^{-k+1}D) \right. \\
 & \quad \left. - \sum_{i=0}^{k-2} \binom{k}{i} (-k+1) \cdots (-2k+2+i) \beta_1(t^{-k+2+i}D^{i+1}, t^{k-1}) \right). \quad (3.10)
 \end{aligned}$$

Taking $k = l + l'$ in (3.7), we obtain (3.5) from (3.7) and (3.10).

Proof of Theorem 2.1. Let β be a 2-cocycle on \mathfrak{a} , then there exists a 2-cocycle β_1 on \mathfrak{a} , β_1 is equivalent to β , and β_1 satisfies (3.2), (3.3). If $\beta_1 = 0$, this implies $\beta_1 = 0\alpha$, $0 \in \mathbb{C}$.

If $\beta_1 \neq 0$, i.e., $\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) \neq 0$ for some $l \geq 0$, $l' \geq 0$, $m \in \mathbb{Z}$. We conclude that $\beta_1(t, t^{-1}) \neq 0$. Otherwise, $\beta_1(t, t^{-1}) = 0$. By Propositions 3.3 and 3.5, $\beta_1(t^m, t^n) = 0$ for $m, n \in \mathbb{Z}$. Now we prove that

$$\beta_1(t^{l+m}D^l, t^{l'+m'}D^{l'}) = 0 \quad (3.11)$$

by induction on $l + l'$. We have known that (3.11) holds for $l + l' = 0$. Assume that (3.11) holds for $l + l' \leq n$. If $l + l' = n + 1$, since the coefficient matrices in (3.4), (3.5) are all nondegenerate and the right hand of (3.4) or (3.5) for $k = n + 1$ is zero, we have

$$\beta_1(t^{-1}D^l, t^{n+2}D^{l'}) = 0,$$

$l + l' = n + 1$. By Propositions 3.3 and 3.5, we see that (3.11) holds for $l + l' = n + 1$. Equation (3.11) implies $\beta_1 = 0$. This is a contradiction. Hence we have $\beta_1(t, t^{-1}) \neq 0$.

Denote $a = \beta_1(t, t^{-1}) \in \mathbb{C}^*$. Define $\beta_2 = a^{-1}\beta_1$, then β_2 is a 2-cocycle and β_2 satisfies (3.2), (3.3). All results for β_1 hold for β_2 and α . Now we prove that $\beta_2 = \alpha$.

By definition, we have $\beta_2(t, t^{-1}) = \alpha(t, t^{-1}) = 1$. From Propositions 3.3 and 3.5, $\beta_2(t^m, t^n) = \alpha(t^m, t^n)$, $m, n \in \mathbb{Z}$. Assume $\beta_2(t^{l+m}D^l, t^{l'+m'}D^{l'}) = \alpha(t^{l+m}D^l, t^{l'+m'}D^{l'})$ for $l + l' \leq n$, if $l + l' = n + 1$. Replace β_1 by β_2 and α , respectively, and set $k = n + 1$; since the coefficient matrices in (3.4), (3.5) are nondegenerate, we have

$$\beta_2(t^{-1}D^l, t^{n+2}D^{l'}) = \alpha(t^{-1}D^l, t^{n+2}D^{l'}),$$

$l + l' = n + 1$. This implies $\beta_2 = \alpha$. Thus we always have $\beta_1 = a\alpha$, $a \in \mathbb{C}$, and β is equivalent to $a\alpha$, $a \in \mathbb{C}$. This completes the proof of the theorem.

4. PROOF OF THEOREM 2.2

Let e_{ij} be the $n \times n$ matrix which is 1 in the i, j -entry and 0 everywhere else, then \mathfrak{A} has a basis [10]

$$\{t^{l+m}D^l \otimes e_{ij} \mid l \geq 0, m \in \mathbb{Z}, 1 \leq i, j \leq n\},$$

and

$$\begin{aligned} & [t^{l+m}D^l \otimes e_{ij}, t^{l'+m'}D^{l'} \otimes e_{i'j'}] \\ &= \delta_{j,i'}(t^{l+m}D^l) \circ (t^{l'+m'}D^{l'}) \\ & \quad \otimes e_{ij'} - \delta_{j',i}(t^{l'+m'}D^{l'}) \circ (t^{l+m}D^l) \otimes e_{ij} \\ &= \delta_{j,i'} \sum_{p=0}^l \binom{l}{p} t^{l+m}(D^{l-p}(t^{l'+m'})) D^{l'+p} \otimes e_{ij'} \\ & \quad - \delta_{j',i} \sum_{q=0}^{l'} \binom{l'}{q} t^{l'+m'}(D^{l'-q}(t^{l+m})) D^{l+q} \otimes e_{i'j}. \end{aligned}$$

In particular,

$$\begin{aligned} [D \otimes e_{ii}, t^{l+m}D^l \otimes e_{ii}] &= [D, t^{l+m}D^l] \otimes e_{ii} \\ &= (l+m) t^{l+m-1}D^l \otimes e_{ii}, \end{aligned} \quad (4.1)$$

$$\begin{aligned} [t^{-1}D^l \otimes e_{ii}, tD \otimes e_{ii}] &= [t^{-1}D^l, tD] \otimes e_{ii} \\ &= (l+1) t^{-1}D^l \otimes e_{ii}, \end{aligned} \quad (4.2)$$

$$[1 \otimes e_{ii}, t^{l+m}D^l \otimes e_{ij}] = t^{l+m}D^l \otimes e_{ij}, \quad i \neq j. \quad (4.3)$$

From the definition (2.5),

$$\begin{aligned} \psi(t^{l+m}D^l \otimes e_{ij}, t^{l'+m'}D^{l'} \otimes e_{i'j'}) &= 0 \\ \Leftrightarrow \delta_{i,j'} \delta_{i',j} \delta_{m,-m'} &= 0 \quad \text{or} \quad \delta_{i,j'} \delta_{i',j} \delta_{m,-m'} \neq 0, \quad 0 \leq m+l \leq l+l' \end{aligned} \quad (4.4)$$

and

$$\psi(t^{l+m}D^l \otimes e_{ij}, t^{l'+m'}D^{l'} \otimes e_{ji}) = \alpha(t^{l+m}D^l, t^{l'+m'}D^{l'}). \quad (4.5)$$

Let φ be a 2-cocycle on \mathfrak{A} , and we define a linear function f_φ on \mathfrak{A} :

$$f_\varphi(t^{l+m-1}D^l \otimes e_{ii}) = \frac{1}{l+m} \varphi(D \otimes e_{ii}, t^{l+m}D^l \otimes e_{ii}), \quad l+m \neq 0,$$

$$f_\varphi(t^{-1}D^l \otimes e_{ii}) = \frac{1}{l+1} \varphi(t^{-1}D^l \otimes e_{ii}, tD \otimes e_{ii}),$$

$$f_\varphi(t^{l+m}D^l \otimes e_{ij}) = \varphi(1 \otimes e_{ii}, t^{l+m}D^l \otimes e_{ij}), \quad i \neq j.$$

Then $\varphi_1 = \varphi - \alpha_{f_\varphi}$ is a 2-cocycle on \mathfrak{A} and φ_1 is equivalent to φ , where α_{f_φ} is the trivial 2-cocycle induced by f_φ . From (4.1), (4.2), (4.3) we have

$$\varphi_1(D \otimes e_{ii}, t^{l+m} D^l \otimes e_{ii}) = 0, \quad l+m \neq 0, \quad (4.6)$$

$$\varphi_1(t^{-1} D^l \otimes e_{ii}, t D \otimes e_{ii}) = 0, \quad (4.7)$$

$$\varphi_1(1 \otimes e_{ii}, t^{l+m} D^l \otimes e_{ij}) = 0, \quad i \neq j. \quad (4.8)$$

Comparing (4.6), (4.7) with (3.2), (3.3), from the proof of Theorem 2.1 we get immediately.

LEMMA 4.1.

$$\begin{aligned} & \varphi_1(t^{l+m} D^l \otimes e_{ii}, t^{l'+m'} D^{l'} \otimes e_{ii}) \\ &= a_i \alpha(t^{l+m} D^l, t^{l'+m'} D^{l'}), \quad a_i \in \mathbb{C}. \end{aligned}$$

LEMMA 4.2. $\varphi_1(t^{l+m} D^l \otimes e_{ii}, t^{l'+m'} D^{l'} \otimes e_{ij}) = 0, i \neq j.$

Proof. First we prove that $\varphi_1(1 \otimes e_{ii}, t^{l+m} D^l \otimes e_{ji}) = 0$ for $i \neq j.$

$$\begin{aligned} & \varphi_1(1 \otimes e_{ii}, t^{l+m} D^l \otimes e_{ji}) \\ &= \varphi_1(1 \otimes e_{ii}, [1 \otimes e_{jj}, t^{l+m} D^l \otimes e_{ji}]) \\ &= \varphi_1([1 \otimes e_{ii}, 1 \otimes e_{jj}], t^{l+m} D^l \otimes e_{ji}) \\ &\quad + \varphi_1(1 \otimes e_{jj}, [1 \otimes e_{ii}, t^{l+m} D^l \otimes e_{ji}]) \\ &= \varphi_1(1 \otimes e_{jj}, -t^{l+m} D^l \otimes e_{ji}). \end{aligned}$$

From (4.8) we have that the last term is zero.

If $i \neq j$, we have

$$\begin{aligned} & \varphi_1(t^{l+m} D^l \otimes e_{ii}, t^{l'+m'} D^{l'} \otimes e_{ij}) \\ &= \varphi_1(t^{l+m} D^l \otimes e_{ii}, [t^{l'+m'} D^{l'} \otimes e_{ij}, 1 \otimes e_{jj}]) \\ &= \varphi_1([t^{l+m} D^l \otimes e_{ii}, t^{l'+m'} D^{l'} \otimes e_{ij}], 1 \otimes e_{jj}) \\ &\quad + \varphi_1(t^{l'+m'} D^{l'} \otimes e_{ij}, [t^{l+m} D^l \otimes e_{ii}, 1 \otimes e_{jj}]) \\ &= \varphi_1((t^{l+m} D^l) \circ (t^{l'+m'} D^{l'}) \otimes e_{ij}, 1 \otimes e_{jj}) \\ &= 0. \end{aligned}$$

LEMMA 4.3. $\varphi_1(f \otimes e_{ij}, g \otimes e_{jk}) = 0, i \neq j, j \neq k, k \neq i, f, g \in \mathfrak{a}.$

Proof.

$$\begin{aligned}
& \varphi_1(f \otimes e_{ij}, g \otimes e_{jk}) \\
&= \varphi_1([1 \otimes e_{ii}, f \otimes e_{ij}], g \otimes e_{jk}) \\
&= \varphi_1([1 \otimes e_{ii}, g \otimes e_{jk}], f \otimes e_{ij}) + \varphi_1(1 \otimes e_{ii}, [f \otimes e_{ij}, g \otimes e_{jk}]) \\
&= \varphi_1(1 \otimes e_{ii}, (f \circ g) \otimes e_{ik}) \\
&= 0 \quad (\text{by Lemma 4.2}).
\end{aligned}$$

LEMMA 4.4. $\varphi_1(f \otimes e_{ij}, g \otimes e_{sk}) = 0$, $i \neq k$, $j \neq s$, $f, g \in \mathfrak{a}$.

Proof. If $i \neq j$,

$$\begin{aligned}
& \varphi_1(f \otimes e_{ij}, g \otimes e_{sk}) \\
&= \varphi_1([1 \otimes e_{ii}, f \otimes e_{ij}], g \otimes e_{sk}) \\
&= \varphi_1(1 \otimes e_{ii}, [f \otimes e_{ij}, g \otimes e_{sk}]) + \varphi_1([1 \otimes e_{ii}, g \otimes e_{sk}], f \otimes e_{ij}) \\
&= \delta_{i,s} \varphi_1(g \otimes e_{ik}, f \otimes e_{ij}) \\
&= -\delta_{i,s} \varphi_1(f \otimes e_{ij}, g \otimes e_{sk}).
\end{aligned}$$

Hence we have $\varphi_1(f \otimes e_{ij}, g \otimes e_{sk}) = 0$.

Since $[\mathfrak{a}, \mathfrak{a}] = \mathfrak{a}$, $\forall f \in \mathfrak{a}$, f can be rewritten as

$$\sum_{r=1}^u [f_{1r}, f_{2r}], \quad f_{1r}, f_{2r} \in \mathfrak{a}.$$

Thus, if $i = j$,

$$\begin{aligned}
& \varphi_1(f \otimes e_{ii}, g \otimes e_{sk}) \\
&= \varphi_1\left(\sum_{r=1}^u [f_{1r} \otimes e_{ii}, f_{2r} \otimes e_{ii}], g \otimes e_{sk}\right) \\
&= \sum_{r=1}^u \varphi_1([f_{1r} \otimes e_{ii}, g \otimes e_{sk}], f_{2r} \otimes e_{ii}) \\
&\quad + \sum_{r=1}^u \varphi_1(f_{1r} \otimes e_{ii}, [f_{2r} \otimes e_{ii}, g \otimes e_{sk}]) \\
&= 0.
\end{aligned}$$

LEMMA 4.5. a_i 's in Lemma 4.1 are equal.

Proof. We prove that if $i \neq j$, $a_i = a_j$. Since

$$\begin{aligned}
 & \varphi_1(t \otimes e_{ii}, t^{-1} \otimes e_{ii}) \\
 &= \varphi_1(t \otimes (e_{ii} - e_{jj}), t^{-1} \otimes e_{ii}) \quad (\text{by Lemma 4.4}) \\
 &= \varphi_1([1 \otimes e_{ij}, t \otimes e_{ji}], t^{-1} \otimes e_{ii}) \\
 &= \varphi_1(1 \otimes e_{ij}, [t \otimes e_{ji}, t^{-1} \otimes e_{ii}]) + \varphi_1([1 \otimes e_{ij}, t^{-1} \otimes e_{ii}], t \otimes e_{ji}) \\
 &= \varphi_1(1 \otimes e_{ij}, 1 \otimes e_{ji}) + \varphi_1(-t^{-1} \otimes e_{ij}, t \otimes e_{ji}), \\
 & \varphi_1(t \otimes e_{jj}, t^{-1} \otimes e_{jj}) \\
 &= \varphi_1(t \otimes e_{jj}, t^{-1} \otimes (e_{jj} - e_{ii})) \quad (\text{by Lemma 4.4}) \\
 &= \varphi_1(t \otimes e_{jj}, [1 \otimes e_{ji}, t^{-1} \otimes e_{ij}]) \\
 &= \varphi_1([t \otimes e_{jj}, 1 \otimes e_{ji}], t^{-1} \otimes e_{ij}) \\
 &\quad + \varphi_1(1 \otimes e_{ji}, [t \otimes e_{jj}, t^{-1} \otimes e_{ij}]) \\
 &= \varphi_1(t \otimes e_{ji}, t^{-1} \otimes e_{ij}) + \varphi_1(1 \otimes e_{ji}, -1 \otimes e_{ij}).
 \end{aligned}$$

This implies $\varphi_1(t \otimes e_{ii}, t^{-1} \otimes e_{ii}) = \varphi_1(t \otimes e_{jj}, t^{-1} \otimes e_{jj})$, that is, $a_i = a_j$.

In order to prove $\varphi_1(f \otimes e_{ii}, g \otimes e_{ii}) = \varphi_1(f \otimes e_{ij}, g \otimes e_{ji})$, $i \neq j$, $f, g \in \mathfrak{a}$, we need the following fact.

LEMMA 4.6. $\varphi_1(1 \otimes e_{ij}, f \otimes e_{ji}) = \varphi_1(f \otimes e_{ij}, 1 \otimes e_{ji})$, $i \neq j$, $f \in \mathfrak{a}$.

Proof.

$$\begin{aligned}
 & \varphi_1(1 \otimes e_{ij}, f \otimes e_{ji}) \\
 &= \varphi_1(1 \otimes e_{ij}, [1 \otimes e_{ji}, f \otimes e_{ii}]) \\
 &= \varphi_1([1 \otimes e_{ij}, 1 \otimes e_{ji}], f \otimes e_{ii}) + \varphi_1(1 \otimes e_{ji}, [1 \otimes e_{ij}, f \otimes e_{ii}]) \\
 &= \varphi_1(1 \otimes e_{ii}, f \otimes e_{ii}) + \varphi_1(1 \otimes e_{ji}, -f \otimes e_{ij}) \quad (\text{by Lemma 4.4}) \\
 &= \varphi_1(f \otimes e_{ij}, 1 \otimes e_{ji}). \quad (\text{by Lemma 4.1}).
 \end{aligned}$$

LEMMA 4.7. $\varphi_1(f \otimes e_{ii}, g \otimes e_{ii}) = \varphi_1(f \otimes e_{ij}, g \otimes e_{ji})$, $i \neq j$, $f, g \in \mathfrak{a}$.

Proof.

$$\begin{aligned}
 & \varphi_1(f \otimes e_{ii}, g \otimes e_{ii}) \\
 &= \varphi_1([f \otimes e_{ij}, 1 \otimes e_{ji}], g \otimes e_{ii}) \quad (\text{by Lemma 4.4}) \\
 &= \varphi_1(f \otimes e_{ij}, [1 \otimes e_{ji}, g \otimes e_{ii}]) \\
 &\quad + \varphi_1([f \otimes e_{ij}, g \otimes e_{ii}], 1 \otimes e_{ji}) \\
 &= \varphi_1(f \otimes e_{ij}, g \otimes e_{ji}) + \varphi_1((-g \circ f) \otimes e_{ij}, 1 \otimes e_{ji}). \quad (4.9)
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 & \varphi_1(f \otimes e_{ii}, g \otimes e_{ii}) \\
 &= \varphi_1(f \otimes e_{jj}, g \otimes e_{jj}) \quad (\text{by Lemma 4.5}) \\
 &= \varphi_1(f \otimes e_{jj}, [g \otimes e_{ji}, 1 \otimes e_{ij}]) \quad (\text{by Lemma 4.4}) \\
 &= \varphi_1([f \otimes e_{jj}, g \otimes e_{ji}], 1 \otimes e_{ij}) \\
 &\quad + \varphi_1(g \otimes e_{ji}, [f \otimes e_{jj}, 1 \otimes e_{ij}]) \\
 &= \varphi_1((f \circ g) \otimes e_{ji}, 1 \otimes e_{ij}) + \varphi_1(f \otimes e_{ij}, g \otimes e_{ji}). \quad (4.10)
 \end{aligned}$$

From (4.9), (4.10), we have

$$\begin{aligned}
 & \varphi_1((f \circ g) \otimes e_{ji}, 1 \otimes e_{ij}) \\
 &= \varphi_1(1 \otimes e_{ji}, (g \circ f) \otimes e_{ij}) \\
 &= \varphi_1((g \circ f) \otimes e_{ji}, 1 \otimes e_{ij}) \quad (\text{by Lemma 4.6}),
 \end{aligned}$$

that is,

$$\varphi_1([f, g] \otimes e_{ji}, 1 \otimes e_{ij}) = 0.$$

Since $[a, a] = a$, we have $\varphi_1(a \otimes e_{ji}, 1 \otimes e_{ij}) = 0$. Thus (4.9) becomes

$$\varphi_1(f \otimes e_{ii}, g \otimes e_{ii}) = \varphi_1(f \otimes e_{ij}, g \otimes e_{ji}).$$

From Lemma 4.1 to Lemma 4.7, we can see easily that $\varphi_1 = a\psi$ for some $a \in \mathbb{C}$. This completes the proof of Theorem 2.2.

ACKNOWLEDGMENTS

I wish to express my sincere thanks to my supervisor Professor Zhe-xian Wan for many helpful suggestions. I am greatly indebted to Professor Victor Kac, whose inspiring lectures at the Graduate School, Academia Sinica, in 1985, led to my research in these problems.

REFERENCES

1. I. B. FRENKEL, Two constructions of affine Lie algebra representations and boson-fermion correspondence in quantum field theory, *J. Funct. Anal.* **44** (1981), 259–327.
2. I. M. GELFAND AND D. B. FUKS, The cohomology of the Lie algebra of the vector fields in a circle, *Funktsional. Anal. i Prilozhen.* **2** (1968), 92–93 [Russian]; *Funct. Anal. Appl.* **2** (1968), 342–343. [English]

3. N. JACOBSON, "Lie Algebras," Interscience, New York, 1962.
4. V. G. KAC, "Infinite Dimensional Lie Algebras," 2nd ed., Combridge Univ. Press, London/New York 1985.
5. V. G. KAC, Lectures on infinite wedge representation and soliton equations, *Shuxue Jinzhan (Chinese Adv. in Math.)* **16** (1987), 343–376.
6. V. G. KAC, A letter to Professor Zhe-xian Wan, 1986.
7. V. G. KAC, D. A. KAZHDAN, J. LEPOWSKY, AND R. L. WILSON, Realization of the basic representation of the Euclidean Lie algebras, *Adv. in Math.* **42** (1981), 83–112.
8. V. G. KAC, AND D. H. PETERSON, Spin and wedge representations of infinite dimensional Lie algebras and groups, *Proc. Nat. Acad. Sci. U.S.A.* **78** (1981), 3308–3312.
9. J. LEPOWSKY AND R. L. WILSON, Construction of the affine Lie algebra $A_1^{(1)}$, *Comm. Math. Phys.* **62** (1978), 43–53.
10. WAN, ZHE-XIAN (WAN, CHÉ-HSIEN), "Lie Algebras," Oxford/Pergamon, London/New York/Elmsford, NY, 1975.